

# HYDROLYSIS REACTIONS

Keyhan BOLELU, Ph.D

## HYDROLYSIS REACTIONS

**HYDROLYSIS** means fragmentation by water.  
(Bond rupture by using water)

Organic molecules that can be fragment by water:  
Carboxylic Acid Derivates (Amides, Acid chlorides, Esters, Acid Anhydrides, Nitriles)



Keyhan BOLELU, Ph.D

## HYDROLYSIS REACTIONS


**Hydrolysis** occurs slowly when it is made only with water, but the reaction rate increases if the reaction is in the presence of acid or base. (Hydrolysis is carried out using an acid or base **catalyser** in aqueous medium.)



The task of the catalysers is to facilitate electron transfer by activating the bond electrons.

Keyhan BOLELU, Ph.D


## HYDROLYSIS OF CARBOXYLIC ACID DERIVATIVES




Keyhan BOLELU, Ph.D

## HYDROLYSIS OF CARBOXYLIC ACID DERIVATIVES

### • HYDROLYSIS in H<sup>+</sup> medium




### • HYDROLYSIS in OH<sup>-</sup> medium





Keyhan BOLELU, Ph.D

## HYDROLYSIS OF ESTERS



Keyhan BOLELU, Ph.D



### Mechanisms of Hydrolysis Reactions of Esters

When hydrolysis is compared with acid and base catalysis, most significant difference can be summarized as follows:

- \* While water, which is a weak nucleophile in acid-catalyzed hydrolysis, is added to carbonyl group (C=O) increasing electrophilic charge by taking proton; OH<sup>-</sup>, which is a strong nucleophile in base-catalyzed hydrolysis, is added to the weak electrophilic C=O bond.
- \* Carboxylic acid salt and alcohol are formed by base-catalyzed hydrolysis.



Keyhan BOLELU, Ph.D

### SAPONIFICATION

Alkaline hydrolysis → SAPONIFICATION

(Soaps are obtained by alkaline hydrolysis of glycerol esters of high-carbon fatty acids.)

Keyhan BOLELU, Ph.D



### HYDROLYSIS OF AMIDE DERIVATIVES

$$\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{NH}_2 \xrightleftharpoons{\text{OH}^-} \text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}^- + \text{NH}_3$$


The appearance of ammonia in the basic environment can be understood from the smell or blueness of the litmus paper.

If the amide used is *N*-substituted amide, primary or secondary amines are formed instead of ammonia.

Keyhan BOLELU, Ph.D

### Mechanisms of Hydrolysis Reactions of Amides


**ACID CATALYSIS**



Keyhan BOLELU, Ph.D

### Mechanisms of Hydrolysis Reactions of Amides

**BASE CATALYSIS**



Keyhan BOLELU, Ph.D

### HYDROLYSIS OF ACID HALIDES

- Since acyl halides and acid anhydrides are active molecules, they hydrolyze under neutral conditions.

$$\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{X}$$

Since the electrophilic charge of carbonyl carbon is greater than the alkyl carbon, the substitution of halogen is easier.

- To avoid hydrolysis of these compounds, acyl halides and anhydrides should be stored under dry  $\text{N}_2$ , the used solvents and reagents should be dry.

Keyhan BOLELU, Ph.D

### HYDROLYSIS OF ACID HALIDES


$$\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{Cl} + \text{H}_2\text{O} \longrightarrow \text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH} + \text{HCl}$$

↓  
pyridine/NaOH (to remove)







Keyhan BOLELU, Ph.D

### HYDROLYSIS OF ACID HALIDES

The hydrolysis reaction of acyl chlorides is nucleophilic acyl substitution.



Keyhan BOLELU, Ph.D



## Oxidation Number (Ox. State)


Ox. step	Primary	Secondary	Tertiary	Quaternary
-4	CH <sub>4</sub>			
-3		RCH <sub>3</sub>		
-2	CH <sub>3</sub> OH		R <sub>2</sub> CH <sub>2</sub>	
-1		RCH <sub>2</sub> OH		R <sub>3</sub> CH
0	CH <sub>2</sub> O		R <sub>2</sub> CHOH	R <sub>3</sub> C
+1		RCHO		R <sub>3</sub> COH
+2	HCOOH		R <sub>2</sub> CO	
+3		RCOOH		
+4	CO <sub>2</sub>			

The oxidation number of a free element is always 0. If an element loses the  $n$  number of electrons, it is considered to be oxidized by  $+n$  value, and if it gains, it is considered to be reduced by  $-n$  value. In organic molecules, the oxidation numbers of the "C" atoms can be calculated by considering each "H",  $-1$ ; each "C",  $0$ ; each heteroatoms,  $+1$  values for the 4 bonds of the "C" atoms.

Keyhan BOLELLU, Ph.D

## Oxidation and Reduction

When an organic compound undergoes reduction, the reduction reactive oxidizes. When an organic compound undergoes oxidation, the oxidation reactive undergoes reduction. Oxidation and reduction reaction take place at the same time.



Keyhan BOLELLU, Ph.D

## Applied in industrial and laboratory Oxidation Methods

- Oxidation in air or with pure oxygen. (These reactions can be accomplished by biologically catalyzing homogeneous or heterogeneous catalysers.)
- Catalytic dehydrogenation at high temperature. (Vanadium oxides)
- Oxidation with some inorganic substances outside oxygen. (The most commonly used reagents: Sodium dichromate + H<sub>2</sub>SO<sub>4</sub> (sulfochromic mixture), in neutral, basic or acidic medium KMnO<sub>4</sub>, concentrated nitric acid, hydrogen peroxide, ozone, some metallic oxide and peroxides, some oxygenated salts.)
- Oxidation with some organic substances or peroxides and peracides.

Keyhan BOLELLU, Ph.D


## Oxidation Reagents

O <sub>2</sub>	HNO <sub>3</sub>	SO <sub>3</sub>	Cl <sub>2</sub>	Ag <sub>2</sub> O	MnO <sub>2</sub>
O <sub>3</sub>	RO-NO	(CH <sub>3</sub> ) <sub>2</sub> S <sup>+</sup> -O <sup>-</sup>	Br <sub>2</sub>	HgO	MnO <sub>4</sub> <sup>-</sup>
H <sub>2</sub> O <sub>2</sub>	Ø-N <sub>2</sub>	SeO <sub>2</sub>	I <sub>2</sub>	Hg(OAc) <sub>2</sub>	CrO <sub>3</sub>
t-BuO-OH	H <sub>2</sub> NCl		NBS	Pb(OAc) <sub>4</sub>	CrO <sub>2</sub> Cl <sub>2</sub>
R-COO-OH	H <sub>3</sub> N <sup>+</sup> -OSO <sub>3</sub> <sup>-</sup>		t-BuOCl	FeCl <sub>3</sub>	OsO <sub>4</sub>
	R <sub>3</sub> N <sup>+</sup> -O <sup>-</sup>			Fe(CN) <sub>6</sub> <sup>-3</sup>	IO <sub>4</sub> <sup>-</sup>

Keyhan BOLELLU, Ph.D

## Dehydrogenation (-2H):


- Heat with Pt, Pd, S or Se
- Substituted quinones (e.g. Chloranil)



Keyhan BOLELLU, Ph.D

## Oxidation of Ethylenic Double Bond and Aromatic Side Chain with KMnO<sub>4</sub>


In dilute basic medium and in the cold (0-5°C), KMnO<sub>4</sub> acts to dihydroxylate the double bond and glycols (1,2-diols) are formed.



Keyhan BOLELLU, Ph.D


## Oxidation of Ethylenic Double Bond and Aromatic Side Chain with $\text{KMnO}_4$

This reaction takes place as cis-addition in stereochemistry.  
For example; cis-cyclohexane-1,2-diol is formed from cyclohexene.




Kayhan BOLELU, Ph.D

The acidic permanganate solution leads to ketone or carboxylic acid formation breaking the double bond according to the environment of the double bond carbon...



Kayhan BOLELU, Ph.D


Oxidation in the side chain takes place in the benzylic carbone. Even if there are alkyl groups including multiple "C" atoms, it is always fragmented to benzoic acid...



Kayhan BOLELU, Ph.D

## Moderate Oxidation with $\text{MnO}_2$


- Freshly prepared  $\text{MnO}_2$  is used in one step oxidation of allylic and benzylic alcohols.
- $\text{MnO}_2 \rightarrow \text{Mn}^{2+}$  is reduced and there is no oxidation to further steps.



Kayhan BOLELU, Ph.D

## Oxidation of the secondary alcohols with chromic acid (Obtaining of Acetone)


- Acetone is obtained by the addition Chromic acid ( $\text{H}_2\text{CrO}_4$ ),  $\text{CrO}_3$  or  $\text{Na}_2\text{Cr}_2\text{O}_7$  to  $\text{H}_2\text{SO}_4$ . While Cr is reduced from +6 to +3, related compound is oxidized.









- Other oxidation reagents including Cr (VI)  
 $\text{Na}_2\text{Cr}_2\text{O}_7$  (sodium bichromate),  $\text{Na}_2\text{CrO}_4$  (sodium chromate),  $\text{H}_2\text{CrO}_4$  (chromic acid),  $\text{CrO}_3$  (chromic oxide),  $\text{CrO}_2\text{Cl}_2$  (chromyl chloride)

Kayhan BOLELU, Ph.D

## Oxidation of ethylenic compounds with $\text{O}_3$ (Ozonolysis)



Kayhan BOLELU, Ph.D



### Preparation of Primary Amines by Action from Amides ve Nitriles

$$\begin{aligned} R-C\equiv N &\xrightarrow{LiAlH_4} R-CH_2-NH_2 \\ R-CONH_2 &\xrightarrow{LiAlH_4} R-CH_2-NH_2 \end{aligned}$$

- LiAlH<sub>4</sub> is converted to alcohols by reducing R-CHO, R<sub>2</sub>C=O, R-COOH, RCH=CH<sub>2</sub>, RCOOR, RCOCl molecules.

$$R-CHO \xrightarrow{LiAlH_4} R-CH_2OH$$

Keyhan BOLELU, Ph.D

### Reaction Mechanism of LiAlH<sub>4</sub>

Keyhan BOLELU, Ph.D

### Reduction of Carbonyl Derivatives with NaBH<sub>4</sub>

NaBH<sub>4</sub> is a weaker reduction agent than LiAlH<sub>4</sub>. It is used to reduce aldehydes and ketones majorly. The reduction process with NaBH<sub>4</sub> can be carried out in an aqueous medium or alcohol.

$$\begin{aligned} H_2C=CH-CO-CH_3 &\xrightarrow{NaBH_4} H_2C=CH-CH(OH)-CH_3 \text{ (3-hydroxy-1-butanone)} \\ R-CHO &\xrightarrow{NaBH_4} R-CH_2OH \end{aligned}$$

Keyhan BOLELU, Ph.D

### Catalytic Reduction (Hydrogenolysis)

Hydrogenolysis is called that is opened by hydrogen and a metal catalyser the bond in the ethylenic and acetylenic unsaturated structures and the bond between carbon and heteroatom. These metals are platinum (Pt), rutenium (Ru), palladium (Pd) and nickel (Ni).

$$\begin{aligned} CH_2=CH_2 &\xrightarrow[Pt \text{ or } Pd]{H_2, \text{ pressure}} CH_3-CH_3 \\ R-C\equiv C-R &\xrightarrow[Pt / CaCO_3]{H_2} R-CH=CH-R \text{ (cis-addition)} \\ R-C(=O)-Cl &\xrightarrow[Pt / H_2SO_4]{H_2} R-CH_2-H + HCl \text{ (Rosemund Reaction)} \\ CH_3-CH_2-CH_2-Cl &\xrightarrow[Pt]{H_2} CH_3-CH_2-CH_3 + HCl \\ R-C(=O)-H &\xrightarrow[NH_3, 150^\circ C]{H_2 / Rh, Ni, \text{ pressure}} R-CH_2-NH_2 \end{aligned}$$

Keyhan BOLELU, Ph.D

### Examples of Reduction using Metal / Acid System

- It is used Zn, Fe, Sn / HCl, H<sub>2</sub>SO<sub>4</sub> and SnCl<sub>2</sub>, FeSO<sub>4</sub>. When metal is treated with acid, the hydrogen gas is formed on the rise.

$$\begin{aligned} Ar-NO_2 &\xrightarrow{Zn / HCl} Ar-NH_2 \\ Ar-C(=O)-R &\xrightarrow[HCl, \text{ heat}]{Zn-Hg} Ar-CH_2-R \\ R-C(=O)-H &\xrightarrow[\text{conc. HCl, heat}]{Zn-Hg} R-CH_3 \end{aligned}$$

- Zn also reduces in aqueous and alcoholic medium.

$$I-CH_2-CH_2-I \xrightarrow[\text{alcohol}]{Zn} CH_2=CH_2$$

Keyhan BOLELU, Ph.D

### Haloform Reaction

- The halogenation of the α-C atom of many ketone compounds can be carried out in the form of total halogenation in a basic medium. So methyl ketones [CH<sub>3</sub>-CO-] contain three halogen atoms in α-C and are formed trihalomethyl ketones.







$$R-C(=O)-CH_3 + X_2 \xrightarrow{\text{basic}} R-COO^- + CHX_3$$







- The haloform reaction takes place via the hypohalogenide derivative of halogens in basic medium.

$$\begin{aligned} X_2 + 2NaOH &\longrightarrow Na^+OX^- + Na^+X^- + H_2O \text{ (hypohalogenite)} \\ Na^+OX^- &\xrightarrow[H_2O]{\text{ketone}} HOX + NaOH \text{ (Hypohalogenous Acid)} \end{aligned}$$

Keyhan BOLELU, Ph.D







## Yield Calculation

$$\% \text{ Yield} = \frac{\text{Practical Yield}}{\text{Theoretical Yield}} \times 100$$

- When the yield calculation is made, it should be determined which of the starting substances is the **Basic Substance**.


Totally spent (consumed) substance by entering all into the reaction

- After the basic substance is detected, the yield calculation is made on this substance. Because, in order to be able to calculate the yield, the starting substance selected must be completely reacted so that mistakes are prevented.

Keyhan BOLELU, Ph.D

## Example 1 (Calculation of yield over weight in aspirin synthesis reaction)

1.3 g salicylic acid and 2 g (1.9 ml) acetic anhydride were reacted to yield 1.6 g aspirin. Calculate the yield of the reaction according to these data.



M.W. = 138.12 g/mole    M.W. = 102.09 g/mole    M.W. = 180.15 g/mole  
 1.3/138.12=0.0094 mole    2/102.09=0.0195 mole    1.6/180.15=0.0089 mole

Keyhan BOLELU, Ph.D

## Solution

All of 0.0094 mol substance react, so salicylic acid is **basic substance**. Yield calculation is made over this substance.

### Calculation of theoretical yield:

From 138.12 g salicylic acid                      180.15 g Aspirin is obtained  
 From 1.3 g salicylic acid                              X g Aspirin is obtained

$$X = 1.3 \times 180.15 / 138.12 = 1.69 \text{ g} = \text{Theoretical yield}$$

The amount obtained in practice = 1.6 g = **Practical yield**

1.69 g product                      %100  
 1.6 g product                        % X


$$X = 1.6 \times 100 / 1.69 = \%94.6$$

**Result : In this reaction, aspirin was synthesized with %94.6 yield.**

Keyhan BOLELU, Ph.D

## Example 2 (Calculation of yield over mole in *p*-aminobenzenesulfonamide synthesis reaction)

191.6 g *p*-aminobenzenesulfonyl chloride and 500 ml ammonia were reacted to yield 135.7 g *p*-aminobenzenesulfonamide. Calculate the yield of the reaction according to these data.



*p*-Aminobenzenesulfonyl chloride    *p*-Aminobenzenesulfonamide  
 M.W. = 191.69 g                      M.W. = 172.21 g

Keyhan BOLELU, Ph.D

## Solution

- p*-Aminobenzenesulfonyl chloride M.W. = 191.69 g
- p*-Aminobenzenesulfonamide    M.W. = 172.21 g

$$\% \text{ Yield} = \frac{\text{Practical Yield}}{\text{Theoretical Yield}} \times 100$$

$$\% 78.84 = \frac{0.7880}{0.9995} \times 100$$

**Result:** In this reaction, *p*-Aminobenzenesulfonamide was synthesized with %78.84 yield.

Keyhan BOLELU, Ph.D